

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**

17
10/6
3

(12) UK Patent Application (19) GB (11) 2 266 529 (13) A

(43) Date of A publication 03.11.1993

(21) Application No 9308684.1

(22) Date of filing 27.04.1993

(30) Priority data

(31) 9209444 (32) 01.05.1992 (33) GB

(71) Applicant

Merck Sharp & Dohme Limited

(Incorporated in the United Kingdom)

Hertford Road, Hoddesdon, Hertfordshire, EN11 9BU,
United Kingdom

(72) Inventors

Walfred S Saari

Graeme I Stevenson

(74) Agent and/or Address for Service

HK Quillin

Merck & Co Inc, European Patent Department, Terlings
Park, Eastwick Road, Harlow, Essex, CM20 2QR,
United Kingdom

(51) INT CL⁵

C07D 217/00, A61K 31/47

(52) UK CL (Edition L)

C2C CAA CBE CWE C1535 C213 C22Y C220 C226
C25Y C250 C251 C31Y C311 C36Y C364 C366
C368 C37X C491 C50Y C500 C624 C628 C65X
C652 C678 C694 C699 C777 C778 C80Y C802
U1S S2416 S2417

(56) Documents cited


EP 0496369 A1 EP 0049605 A1
Synth. Commun., 22(19), 2745-58
Helv. Chim. Acta., (7), 1944-54
J. Med. Chem., 29(10), 1953-61

(58) Field of search

UK CL (Edition L) C2C
INT CL⁵ C07D
Online databases: CAS ONLINE

(54) Tetrahydroisoquinoline derivatives

(57) Compounds of formula (I), and salts and prodrugs thereof



(I)

wherein:

R¹ is H, C₁₋₆alkyl or optionally substituted phenyl;

R² is H, C₁₋₆alkyl, COR⁷, COOR⁷, CONHR⁷ or optionally substituted phenyl; C₁₋₆alkyl; R⁷ is C₁₋₆alkyl or optionally substituted phenyl;

R³ is optionally substituted phenyl;

each R⁴ is C₁₋₆alkyl, C₁₋₆alkoxy, halo or trifluoromethyl;

R⁵ is H or C₁₋₆alkyl;

R⁶ is H, C₁₋₆alkyl or optionally substituted phenyl;

n is 0, 1, 2, 3 or 4;

X is CH₂ or C=O;

Y is O, S, CH₂ or NR¹⁰; where R¹⁰ is H, C₁₋₆alkyl or COR⁸; or X and Y together represent CH=CH; with the proviso that X is not C=O when Y is CH₂;

are tachykinin receptor antagonists useful in therapy.

GB 2 266 529 A

- 1 -

TETRAHYDROISOQUINOLINE DERIVATIVES

5 This invention relates to a class of compounds which are useful as tachykinin antagonists. More particularly, the compounds of the invention are tetrahydroisoquinoline derivatives.

10 The tachykinins are a group of naturally occurring peptides found widely distributed throughout mammalian tissues, both within the central nervous system and in peripheral nervous and circulatory systems.

The structures of three known mammalian tachykinins are as follows:

Substance P:

15 Arg-Pro-Lys-Pro-Gln-Gln-Phe-Phe-Gly-Leu-Met-NH₂

Neurokinin A:

His-Lys-Thr-Asp-Ser-Phe-Val-Gly-Leu-Met-NH₂

Neurokinin B:

Asp-Met-His-Asp-Phe-Phe-Val-Gly-Leu-Met-NH₂

20 For example, substance P is believed inter alia to be involved in the neurotransmission of pain sensations [Otsuka et al, "Role of Substance P as a Sensory Transmitter in Spinal Cord and Sympathetic Ganglia" in 1982 Substance P in the Nervous System, Ciba Foundation Symposium 91, 13-34 (published by Pitman) and Otsuka and Yanagisawa, "Does Substance P Act as a Pain Transmitter?" TIPS (Dec. 1987) 8 506-510], specifically in the transmission of pain in migraine (B.E.B. Sandberg et al, J. Med Chem, (1982) 25 1009; S.L. Shephard et
25 al., Br. J. Pharmacol. (1993), 108, 11-12) and in arthritis [Levine et al in Science (1984) 226 547-549]. These peptides have also been implicated in gastrointestinal (GI) disorders and diseases of the GI tract such as inflammatory bowel disease [Mantyh et al in

Neuroscience (1988) 25 (3) 817-37 and D. Regoli in "Trends in Cluster Headache" Ed. Sicuteri et al Elsevier Scientific Publishers, Amsterdam (1987) page 85)]. It is also hypothesised that there is a neurogenic mechanism
5 for arthritis in which substance P may play a role [Kidd et al "A Neurogenic Mechanism for Symmetrical Arthritis" in The Lancet, 11 November 1989 and Grönblad et al "Neuropeptides in Synovium of Patients with Rheumatoid Arthritis and Osteoarthritis" in J. Rheumatol. (1988)
10 15(12) 1807-10]. Therefore, substance P is believed to be involved in the inflammatory response in diseases such as rheumatoid arthritis and osteoarthritis [O'Byrne et al in Arthritis and Rheumatism (1990) 33 1023-8]. Other disease areas where tachykinin antagonists are believed
15 to be useful are allergic conditions [Hamelet et al Can. J. Pharmacol. Physiol. (1988) 66 1361-7], immunoregulation [Lotz et al Science (1988) 241 1218-21 and Kimball et al, J. Immunol. (1988) 141 (10) 3564-9] vasodilation, bronchospasm, reflex or neuronal control of
20 the viscera [Mantyh et al, PNAS (1988) 85 3235-9] and, possibly by arresting or slowing β -amyloid-mediated neurodegenerative changes [Yankner et al Science (1990) 250, 279-82] in senile dementia of the Alzheimer type, Alzheimer's disease and Down's Syndrome.

25 Substance P may also play a role in demyelinating diseases such as multiple sclerosis and amyotrophic lateral sclerosis [J. Lubner-Narod et. al., poster presented at C.I.N.P. XVIIIth Congress, 28th June-2nd July, 1992], and in disorders of bladder function
30 such as bladder detrusor hyper-reflexia (Lancet, 16th May, 1992, 1239).


It has furthermore been suggested that tachykinins have utility in the following disorders: depression, dysthymic disorders, chronic obstructive

airways disease, hypersensitivity disorders such as
poison ivy, vasospastic diseases such as angina and
Reynauld's disease, fibrosing and collagen diseases such
as scleroderma and eosinophillic fascioliasis, reflex
5 sympathetic dystrophy such as shoulder/hand syndrome,
addiction disorders such as alcoholism, stress related
somatic disorders, neuropathy, neuralgia, disorders
related to immune enhancement or suppression such as
systemic lupus erythmatosis (European patent application
10 no. 0 436 334), ophthalmic disease such as conjunctivitis,
vernal conjunctivitis, and the like, and cutaneous
diseases such as contact dermatitis, atropic dermatitis,
urticaria, and other eczematoid dermatitis (European
patent application no. 0 394 989) and emesis (European
15 patent application no. 0 533 280).

In view of their metabolic instability, peptide
derivatives are likely to be of limited utility as
therapeutic agents. It is for this reason that non-
peptide tachykinin antagonists are sought.

20 Tetrahydroisoquinoline derivatives said to be
useful as substance P antagonists are disclosed in WO
92/06079. There is no disclosure of the substitution
pattern of the compounds of the present invention.

The present invention provides a compound of
25 formula (I), or a salt or prodrug thereof:



(I)

wherein:

5 R^1 represents H, C_{1-6} alkyl or phenyl optionally substituted by 1, 2 or 3 groups selected from C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, halo, cyano, nitro, trifluoromethyl, trimethylsilyl, $-OR^a$, SR^a , SOR^a , SO_2R^a , $-NR^aR^b$, $-NR^aCOR^b$, $-NR^aCO_2R^b$, $-CO_2R^a$ or $-CONR^aR^b$;

R^2 represents H, C_{1-6} alkyl, COR^7 , $COOR^7$, $CONHR^7$ or phenyl(C_{1-4} alkyl) optionally substituted in the phenyl ring by C_{1-6} alkyl, C_{1-6} alkoxy, halo or trifluoromethyl;

10 R^3 represents phenyl optionally substituted by 1, 2 or 3 groups selected from C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, halo, cyano, nitro, trifluoromethyl, trimethylsilyl, $-OR^a$, SR^a , SOR^a , SO_2R^a , $-NR^aR^b$, $-NR^aCOR^b$, $-NR^aCO_2R^b$, $-CO_2R^a$ and $-CONR^aR^b$;

15 each R^4 independently represents C_{1-6} alkyl, C_{1-6} alkoxy, halo or trifluoromethyl;

R^5 represents H or C_{1-6} alkyl;

20 R^6 represents H, C_{1-6} alkyl or phenyl optionally substituted by 1, 2 or 3 groups selected from C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, halo, cyano, nitro, trifluoromethyl, trimethylsilyl, $-OR^a$, SR^a , SOR^a , SO_2R^a , $-NR^aR^b$, $-NR^aCOR^b$, $-NR^aCO_2R^b$, $-CO_2R^a$ and $-CONR^aR^b$;

R^7 is C_{1-6} alkyl or phenyl optionally substituted by C_{1-6} alkyl, C_{1-6} alkoxy, halo or trifluoromethyl;

25 n is 0, 1, 2, 3 or 4;

X represents CH_2 or $C=O$;

Y represents O, S, CH_2 , or NR^{10} where R^{10} represents H, C_{1-6} alkyl or COR^7 ; or X and Y together represent $CH=CH$; with the proviso that X is not $C=O$ when

30 Y is CH_2 ; and

R^a and R^b each independently represent H, C_{1-6} alkyl, trifluoromethyl or phenyl optionally substituted by C_{1-6} alkyl, halo or trifluoromethyl.

As used herein, the definition of each expression, when it occurs more than once in any structure, is intended to be independent of its definition elsewhere in the same structure.

5 The alkyl, alkenyl and alkynyl groups referred to with respect to the above formula may represent straight, branched or cyclic groups or combinations thereof. Thus, for example, suitable alkyl groups include methyl, ethyl, n- or iso-propyl, n-, sec-, iso-
10 or tert-butyl, cyclopropyl, cyclobutyl, cyclopentyl or cyclohexyl, and cycloalkyl-alkyl groups such as cyclopropylmethyl; suitable alkenyl groups include vinyl and allyl; and suitable alkynyl groups include propargyl.

15 The term "halo" as used herein includes fluoro, chloro, bromo and iodo.

 Suitably, R^1 represents unsubstituted or substituted phenyl, preferably unsubstituted phenyl.

 Suitably R^2 represents H or C_{1-6} alkyl, preferably H or methyl.


20 Suitable values for R^4 include, for example, methyl, methoxy, chloro, fluoro and trifluoromethyl.

 Preferably R^5 and R^6 are both H.

 Preferably n is 0

25 Preferably Y represents O or NR^{10} , more preferably O.

 A particular sub-class of compounds according to the invention is represented by the compounds of formula (IIa), and salts and prodrugs thereof:



(IIa)

wherein:


10 R^2 , X and Y are each as defined above with reference to formula (I); and R^{21} and R^{22} each independently represent H, C_1 -6alkyl, halo, cyano, nitro, trifluoromethyl, hydroxy, C_1 -6alkoxy, phenoxy or amino.

15 Preferred are compounds of formula (IIa) wherein R^{21} and R^{22} are selected from H, methyl, ethyl, t-butyl, methoxy, fluoro, chloro, bromo, iodo and trifluoromethyl. Preferably R^{21} and R^{22} are both other than H, more preferably C_1 -6alkyl, halo or
20 trifluoromethyl, and are located at the 3- and 5-positions of the phenyl ring. Compounds wherein both R^{21} and R^{22} represent trifluoromethyl are particularly preferred.

25 A preferred sub-class of compounds according to the invention are those wherein Y represents O.

It is preferred that the substituents R^1 and X- are in the trans orientation, that is to say, the relative stereochemistry indicated in formula (Ia) is preferred:

30



(Ia)

10 For use in medicine, the salts of the compounds
of formula (I) will be pharmaceutically acceptable salts.
Other salts may, however, be useful in the preparation of
the compounds according to the invention or of their
pharmaceutically acceptable salts. Suitable
15 pharmaceutically acceptable salts of the compounds of
this invention include acid addition salts which may, for
example, be formed by mixing a solution of the compound
according to the invention with a solution of a
pharmaceutically acceptable acid such as hydrochloric
20 acid, sulphuric acid, oxalic acid, fumaric acid, p-
toluenesulphonic acid, maleic acid, succinic acid, acetic
acid, citric acid, tartaric acid, carbonic acid or
phosphoric acid. Salts of amine groups may also comprise
quaternary ammonium salts in which the amino nitrogen
25 atom carries a suitable organic group such as an alkyl,
alkenyl, alkynyl or aralkyl moiety. Furthermore, where
the compounds of the invention carry an acidic moiety,
suitable pharmaceutically acceptable salts thereof may
include metal salts such as alkali metal salts, e.g.
30 sodium or potassium salts; and alkaline earth metal
salts, e.g. calcium or magnesium salts.

The present invention includes within its scope
prodrugs of the compounds of formula (I) above. In
general, such prodrugs will be functional derivatives of

the compounds of formula (I) which are readily convertible in vivo into the required compound of formula (I). Conventional procedures for the selection and preparation of suitable prodrug derivatives are described, for example, in "Design of Prodrugs", ed. H. Bundgaard, Elsevier, 1985.

The compounds according to the invention have at least one asymmetric centre, and may accordingly exist both as enantiomers and as diastereoisomers. It is to be understood that all such isomers and mixtures thereof are encompassed within the scope of the present invention.

The invention also provides pharmaceutical compositions comprising one or more compounds of this invention in association with a pharmaceutically acceptable carrier. Preferably these compositions are in unit dosage forms such as tablets, pills, capsules, powders, granules, solutions or suspensions, or suppositories, for oral, parenteral or rectal administration, or administration by inhalation or insufflation.

The invention further provides a process for the preparation of a pharmaceutical composition comprising a compound of formula (I), or a salt or prodrug thereof, and a pharmaceutically acceptable carrier, which process comprises bringing a compound of formula (I), or a salt or prodrug thereof into association with a pharmaceutically acceptable carrier.

For preparing solid compositions such as tablets, the principal active ingredient is mixed with a pharmaceutical carrier, e.g. conventional tableting ingredients such as corn starch, lactose, sucrose, sorbitol, talc, stearic acid, magnesium stearate, dicalcium phosphate or gums, and other

pharmaceutical diluents, e.g. water, to form a solid preformulation composition containing a homogeneous mixture of a compound of the present invention, or a non-toxic pharmaceutically acceptable salt thereof. When referring to these preformulation compositions as homogeneous, it is meant that the active ingredient is dispersed evenly throughout the composition so that the composition may be readily subdivided into equally effective unit dosage forms such as tablets, pills and capsules. This solid preformulation composition is then subdivided into unit dosage forms of the type described above containing from 0.1 to about 500 mg of the active ingredient of the present invention. The tablets or pills of the novel composition can be coated or otherwise compounded to provide a dosage form affording the advantage of prolonged action. For example, the tablet or pill can comprise an inner dosage and an outer dosage component, the latter being in the form of an envelope over the former. The two components can be separated by an enteric layer which serves to resist disintegration in the stomach and permits the inner component to pass intact into the duodenum or to be delayed in release. A variety of materials can be used for such enteric layers or coatings, such materials including a number of polymeric acids and mixtures of polymeric acids with such materials as shellac, cetyl alcohol and cellulose acetate.

The liquid forms in which the novel compositions of the present invention may be incorporated for administration orally or by injection include aqueous solutions, suitably flavoured syrups, aqueous or oil suspensions, and flavoured emulsions with edible oils such as cottonseed oil, sesame oil, coconut oil or peanut oil, as well as elixirs and similar pharmaceutical

vehicles. Suitable dispersing or suspending agents for aqueous suspensions include synthetic and natural gums such as tragacanth, acacia, alginate, dextran, sodium carboxymethylcellulose, methylcellulose, polyvinylpyrrolidone or gelatin.

Compositions for inhalation or insufflation include solutions and suspensions in pharmaceutically acceptable, aqueous or organic solvents, or mixtures thereof, and powders. The liquid or solid compositions may contain suitable pharmaceutically acceptable excipients as set out above. Preferably the compositions are administered by the oral or nasal respiratory route for local or systemic effect. Compositions in preferably sterile pharmaceutically acceptable solvents may be nebulised by use of inert gases. Nebulised solutions may be breathed directly from the nebulising device or the nebulising device may be attached to a face mask, tent or intermittent positive pressure breathing machine. Solution, suspension or powder compositions may be administered, preferably orally or nasally, from devices which deliver the formulation in an appropriate manner.

The substance P antagonising activity of the compounds described herein was evaluated using the human NK1R assay described in published European patent application no. 0 528 495. The method essentially involves determining the concentration of the test compound required to reduce by 50% the amount of radiolabelled substance P binding to human NK1R, thereby affording an IC₅₀ value for the test compound. The compounds of Examples 1-3 were found to have IC₅₀ values less than 1 μ M.

The compounds of formula (I) are of value in the treatment of a wide variety of clinical conditions which are characterised by the presence of an excess of

tachykinin, in particular substance P, activity. These may include disorders of the central nervous system such as anxiety, depression, psychosis and schizophrenia; neurodegenerative disorders such as dementia, including
5 senile dementia of the Alzheimer type, Alzheimer's disease and Down's syndrome; demyelinating diseases such as multiple sclerosis (MS) and amyotrophic lateral sclerosis (ALS; Lou Gehrig's disease) and other neuropathological disorders such as peripheral
10 neuropathy, for example, diabetic or chemotherapy-induced neuropathy, and postherpetic and other neuralgias; respiratory diseases such as chronic obstructive airways disease, bronchopneumonia, bronchospasm and asthma; inflammatory diseases such as inflammatory bowel disease,
15 psoriasis, fibrositis, osteoarthritis and rheumatoid arthritis; allergies such as eczema and rhinitis; hypersensitivity disorders such as poison ivy; ophthalmic diseases such as conjunctivitis, vernal conjunctivitis, and the like; cutaneous diseases such as contact
20 dermatitis, atopic dermatitis, urticaria, and other eczematoid dermatitis; addiction disorders such as alcoholism; stress related somatic disorders; reflex sympathetic dystrophy such as shoulder/hand syndrome; dysthymic disorders; adverse immunological reactions such
25 as rejection of transplanted tissues and disorders related to immune enhancement or suppression such as systemic lupus erythematosus; gastrointestinal (GI) disorders and diseases of the GI tract such as disorders associated with the neuronal control of viscera such as
30 ulcerative colitis, Crohn's disease and incontinence; emesis, including acute, delayed and anticipatory emesis, for example, induced by chemotherapy, radiation, toxins, pregnancy, vestibular disorders, surgery, migraine and variations in intracranial pressure; disorders of bladder

function such as bladder detrusor hyper-reflexia;
fibrosing and collagen diseases such as scleroderma and
eosinophilic fascioliasis; disorders of blood flow caused
by vasodilation and vasospastic diseases such as angina,
5 migraine and Reynaud's disease; and pain or nociception,
for example, that attributable to or associated with any
of the foregoing conditions, especially the transmission
of pain in migraine. For example, the compounds of
formula (I) may suitably be used in the treatment of
10 disorders of the central nervous system such as anxiety,
psychosis and schizophrenia; neurodegenerative disorders
such as senile dementia of the Alzheimer type,
Alzheimer's disease and Down's syndrome; respiratory
diseases, particularly those associated with excess mucus
15 secretion, such as chronic obstructive airways disease,
bronchopneumonia, chronic bronchitis, cystic fibrosis and
asthma, and bronchospasm; inflammatory diseases such as
inflammatory bowel disease, osteoarthritis and rheumatoid
arthritis; adverse immunological reactions such as
20 rejection of transplanted tissues; gastrointestinal (GI)
disorders and diseases of the GI tract such as disorders
associated with the neuronal control of viscera such as
ulcerative colitis, Crohn's disease and incontinence;
disorders of blood flow caused by vasodilation; and pain
25 or nociception, for example, that attributable to or
associated with any of the foregoing conditions or the
transmission of pain in migraine.

The compounds of formula (I) are particularly
useful in the treatment of pain or nociception and/or
30 inflammation and disorders associated therewith such as,
for example, neuropathy, such as diabetic and
chemotherapy-induced neuropathy, postherpetic and other
neuralgias, asthma, osteoarthritis, rheumatoid arthritis
and especially migraine.

The present invention further provides a compound of formula (I) for use in therapy. According to a further or alternative aspect, the present invention provides a compound of formula (I) for use in the
5 manufacture of a medicament for the treatment of physiological disorders associated with an excess of tachykinins, especially substance P. The present invention also provides a method for the the treatment or prevention of physiological disorders associated with an
10 excess of tachykinins, especially substance P, which method comprises administration to a patient in need thereof of a tachykinin reducing amount of a compound of formula (I) or a composition comprising a compound of formula (I).

15 For the treatment of certain conditions it may be desirable to employ a compound according to the present invention in conjunction with another pharmacologically active agent. For example, for the treatment of respiratory diseases such as asthma, a
20 compound of formula (I) may be used in conjunction with a bronchodilator, such as a β_2 -adrenergic receptor antagonist or tachykinin antagonist which acts at NK-2 receptors. The compound of formula (I) and the bronchodilator may be administered to a patient
25 simultaneously, sequentially or in combination.

The present invention accordingly provides a method for the treatment of a respiratory disease, such as asthma, which method comprises administration to a patient in need thereof of an effective amount of a
30 compound of formula (I) and an effective amount of a bronchodilator.

The present invention also provides a composition comprising a compound of formula (I), a

In the treatment of the conditions associated with an excess of tachykinins, a suitable dosage level is about 0.001 to 50 mg/kg per day, in particular about 0.01 to about 25 mg/kg, such as from about 0.05 to about 10 mg/kg per day. For example, in the treatment of conditions involving the neurotransmission of pain sensations, a suitable dosage level is about 0.001 to 25 mg/kg per day, preferably about 0.005 to 10 mg/kg per day, and especially about 0.005 to 5 mg/kg per day. The compounds may be administered on a regimen of 1 to 4 times per day, preferably once or twice per day.

*c1ccc2c(c1)c(c[nH]2)C(R1)C(X)Y

(1 1 1)

The reaction is conveniently carried out in a suitable organic solvent, such as an ether, for example, tetrahydrofuran.

Suitable bases of use in the reaction include alkali or alkaline earth metal hydrides, for example, sodium hydride, and alkali metal carbonates, such as caesium carbonate.


5 The compounds of the invention wherein Y is a group NR^{10} and X represents C=O may be prepared from the compounds of formula (III) wherein Y is O and X represents C=O by reaction with a compound of formula $\text{R}^3\text{CH}_2\text{NHR}^{10}$, where R^3 and R^{10} are as defined for formula
10 (I), in the presence of a coupling agent, such as dicyclohexylcarbodiimide.

The reaction is suitably effected in an aprotic organic solvent, such as dichloromethane or dimethylformamide, or a mixture thereof.

15 The compounds according to the invention wherein Y is NR^{10} and X is CH_2 may be prepared from the corresponding compounds of formula (I) wherein X is C=O by reduction.

20 Suitable reducing agents of use in the reaction include borane and metal hydrides, such as lithium aluminium hydride. The reaction is conveniently effected in a suitable organic solvent, such as an ether, for example, tetrahydrofuran.

25 Compounds of formula (I) wherein X and Y together represent CH=CH may be prepared from intermediates of formula (IV)



(IV)


$$R^{11} \quad {}_3P \quad \begin{array}{c} \diagup \\ \diagdown \end{array} \begin{array}{c} R^3 \\ R^5 \\ R^6 \end{array}$$

(v)

Suitable bases include the alkali or alkaline earth metal salts of alcohols, such as, for example potassium t-butoxide.

Compounds of formula (I) wherein X and Y both represent groups CH_2 may be prepared from the corresponding compounds of formula (I) wherein X and Y together represent $\text{CH}=\text{CH}$, by reduction. Suitable procedures and reagents will be readily apparent to persons skilled in the art. For example, the conversion may be achieved using catalytic hydrogenation or dissolving metal reduction with, for example, magnesium in methanol.

Compounds of formula (III) wherein X is C=O and Y is O may be prepared from intermediates of formula (VI)



(VI)

wherein R^1 , R^2 , R^4 and n are as defined for formula (I),
 10 by reaction with formaldehyde in the presence of a
 mineral acid, such as hydrochloric acid.

Conveniently, the reaction is effected in
 aqueous solution.


Compounds of formula (III) wherein X is CH_2 may
 15 be prepared from the corresponding compounds of formula
 (III) wherein X is $C=O$, by reduction, for example, using
 a metal hydride, such as lithium aluminium hydride.

Compounds of formula (III) wherein Y is S may
 be prepared from the corresponding compounds wherein Y is
 20 O by treating the latter compounds with Lawesson's
 reagent or phosphorus pentasulphide in a suitable
 solvent, e.g. pyridine, at ambient or elevated
 temperature, suitably at the reflux temperature of the
 chosen solvent.

Intermediates of formula (IV) may be prepared
 25 from intermediates of formula (III) wherein X is $C=O$ and
 Y is O by reduction, for example using a metal hydride
 reducing agent, such as diisobutylaluminium hydride.

Intermediates of formula (V) are commercially
 30 available or can be prepared from the corresponding
 halides, for example, using the Arbuzov reaction.

Intermediates of formula (VI) may be prepared
 from compounds of formula (VII)




(VII)

wherein R^1 , R^4 and n are as defined for formula (I) and
 10 Ph represents phenyl, by hydrolysis.

The reaction is conveniently effected by heating a solution of the compound of formula (VII) in concentrated hydrochloric acid at reflux.


Intermediates of formula (VII) may be prepared
 15 from the commercially available compound of formula (VIII)



(VIII)

by reaction with compounds of formula (IX)

25



(IX)

wherein R^1 , R^4 and n are as defined for formula (I) and Hal is halo, such as chloro, bromo or iodo, in the presence of a base.

Suitable bases of use in the reaction include metal hydroxides, for example, sodium hydroxide. The reaction is conveniently effected in a mixture of water and a suitable organic solvent, such as a hydrocarbon, for example, toluene, in the presence of a phase transfer catalyst, such as benzyltrimethyl ammonium chloride.

Compounds of formula (IX) may be prepared according to the procedure described by E. J. Corey, Tetrahedron Lett., 1972, 4339, or methods analogous thereto.

Where the above-described process for the preparation of the compounds according to the invention gives rise to mixtures of stereoisomers these isomers may, if desired, be separated, suitably by conventional techniques such as preparative chromatography.

The novel compounds may be prepared in racemic form, or individual enantiomers may be prepared either by enantiospecific synthesis or by resolution. The novel compounds may, for example, be resolved into their component enantiomers by standard techniques, such as the formation of diastereomeric pairs by salt formation with an optically active acid, such as (-)-di-p-toluoyl-d-tartaric acid and/or (+)-di-p-toluoyl-l-tartaric acid followed by fractional crystallization and regeneration of the free base. The novel compounds may also be resolved by formation of diastereomeric esters or amides, followed by chromatographic separation and removal of the chiral auxiliary.

During any of the above synthetic sequences it may be necessary and/or desirable to protect sensitive or reactive groups on any of the molecules concerned. This

may be achieved by means of conventional protecting groups, such as those described in Protective Groups in Organic Chemistry, ed. J.F.W. McOmie, Plenum Press, 1973; and T.W. Greene and P.G.M. Wuts, Protective Groups in Organic Synthesis, John Wiley & Sons, 1991. The
5 protecting groups may be removed at a convenient subsequent stage using methods known from the art.

The following Examples illustrate the preparation of compounds according to the invention.

EXAMPLE 1

cis 4-Phenyl-3-(3,5-bis[trifluoromethyl]-1,2,3,4-tetrahydroisoquinoline Hydrochloride

5 a) Diphenylalaninol

Diphenylalanine methyl ester (6.3g) in dry tetrahydrofuran was added dropwise to a stirred solution of lithium aluminum hydride (986mg) in dry tetrahydrofuran. Once addition was complete the solution was warmed to reflux for one hour and then cooled to room temperature. Water (10ml), 2N sodium hydroxide solution (10ml), and a further aliquot of water (10ml) were added and the solution stirred at room temperature for one hour. The reaction mixture was then filtered through hyflo, diluted with water (100ml) and extracted into ethyl acetate. The organic layers were separated, dried (MgSO₄), filtered and the solvent removed under reduced pressure, to afford the product as a clear oil (5.02g). ¹H NMR (360MHz, CDCl₃) δ 3.30 (1H, dd, J = 8.0, 3.0Hz, CHHOH), 3.53 (1H, dd, J = 8.0, 1.0Hz, CHH-OH), 3.62 (1H, m, CH-CH₂OH), 3.75 (1H, d, J = 7.0Hz, CHCHCH₂OH), 6.9-7.3 (10H, m, ArH). m/z (CI⁺) 228.

25 b) N-t-Butoxycarbonyl diphenylalaninol

Di-t-butyl dicarbonate (5.23g) was added to a stirred solution of diphenylalaninol (5.02g) in dichloromethane. The solution was stirred for eighteen hours at room temperature. After this time the solvent was removed under reduced pressure to afford a yellow oil. Recrystallisation from hexane afforded the pure product as yellow needles (3.83g). mp 78-79°C. ¹H NMR

(360MHz, CDCl_3) δ 1.34 (9H, s, $\text{C}(\text{CH}_3)_3$), 3.47 (1H, m, CHH-OH), 3.67 (1H, m, CHH-OH), 4.13 (1H, d, $J = 7.0\text{Hz}$, CHCHCH_2OH), 4.46 (1H, m, CHCHCH_2OH), 4.59 (1H, brs, NH), 7.13-7.35 (10H, m, ArH); m/z (Cl^+) 328.

5

c) N-t-Butyloxycarbonyl O-[3,5-bis(trifluoromethyl)benzyl] diphenylalaninol

Sodium hydride (288mg, 80%) was added to an ice cold solution of N-t-butoxycarbonyl diphenylalaninol (3.83g) and 3,5-bis(trifluoromethyl)benzyl bromide (3.59g) in dry dimethylformamide. The resulting mixture was allowed to warm to room temperature overnight. The reaction mixture was poured into water and extracted into ethyl acetate. The organic layers were separated, dried (MgSO_4), filtered and the solvent removed under reduced pressure. Product isolated by flash chromatography as an oil (2.4g). ^1H NMR (360MHz, CDCl_3) 1.3 (9H, s, $\text{C}(\text{CH}_3)_3$), 3.3 (1H, d, $J = 8.0\text{Hz}$, CHCHH-O), 3.45 (1H, d, $J = 8.0\text{Hz}$, CHCHH-O-), 4.2 (1H, m, $\text{CHCH}_2\text{-O-}$), 4.4-4.6 (3H, m, CHCH and $\text{Ar-CH}_2\text{-O-CH}_2\text{CH}$), 7.1-7.3 (10H, m, ArH), 7.7 (2H, s, $\text{CF}_3\text{C-CH-C}$), 7.8 (1H, s, $\text{CF}_3\text{C-CH-CCF}_3$); m/z (Cl^+) 553.

20

d) O-(3,5-Bistrifluoromethyl)benzyl diphenylalaninol

Trifluoroacetic acid (25ml) was added to a stirred solution of N-t-butyloxycarbonyl O-[3,5-(bistrifluoromethyl)benzyl] diphenylalaninol (3.49g) in dry dichloromethane. The resulting solution was stirred for two hours at room temperature. The reaction mixture was then partitioned between dichloromethane and saturated sodium carbonate solution. The organic layer was separated, washed with water, dried (MgSO_4), filtered and the solvent removed to afford the product as an oil (1.8g). ^1H NMR (360MHz, CDCl_3) 3.2 (1H, d, $J = 8.0\text{Hz}$, CHCHH-O-), 3.5 (1H, d,

25

30

$J = 8.0\text{Hz}$, CHCHH-O), 4.15 (1H, m, $\text{CHCH}_2\text{-O-}$), 4.4-4.6 (3H, m, CHH and $\text{ArCH}_2\text{-OCH}_2\text{CH}$), 7.0-7.3 (10H, m, ArH), 7.61 (2H, s, CF_3CCHC), 7.68 (1H, s, $\text{CF}_3\text{CCHCCF}_3$). m/z (CI^+) 453.

5 e) N-Formyl O-[3,5-bis(trifluoromethyl)benzyl diphenylalaninol

1-(3-dimethylaminopropyl)-3-ethyl carbodiimide hydrochloride (1.72g) was added to a stirred solution of
 10 O-[3,5-bis(trifluoromethyl)benzyl diphenylalaninol (2.54g) and formic acid (7.5ml) in dry dichloromethane. The resulting mixture was stirred at room temperature for six hours and then partitioned between water and dichloromethane. The organic layers were separated, washed with citric acid and water, dried
 15 (MgSO_4), filtered and solvent removed. Re-crystallisation from ethyl acetate/hexane afforded the product was yellow needles (2.6g). ^1H NMR (360MHz, CDCl_3) 3.35 (1H, dd, $J = 8.0, 1.0\text{Hz}$, $\text{CHH O-CH}_2\text{Ar}$), 3.52 (1H, dd, $J = 8.0, 1.0\text{Hz}$, CHHOCH_2Ar), 4.4 (1H, d, $J = 9.0\text{Hz}$, CHH-Ar), 4.51 (1H, d, $J = 4.0\text{Hz}$,
 20 $\text{CH-CHCH}_2\text{O}$), 4.58 (1H, d, $J = 9.0\text{Hz}$, CHH-Ar), 5.07 (1H, m, $\text{CH-CH}_2\text{OCH}_2$), 5.71 (1H, brd, NH), 7.2-7.3 (10H, m, ArH), 7.7 (2H, s, CCH-CCF_3), 7.79 (1H, s, $\text{CF}_3\text{C-CH-CCF}_3$), 8.05 (1H, s, CHO). m/z (CI^-) 481.

25 f) 4-Phenyl. 3-(3,5-bis[trifluoromethyl]benzyloxymethyl 1,2,3,4-tetrahydroisoquinoline hydrochloride

Oxalyl chloride (513mg) was added to a solution of
 30 N-formyl O-(3,5-bis[trifluoromethyl]benzyl) diphenylalaninol (1.62g) in dry dichloromethane at room temperature. The resulting solution was stirred for one hour at room temperature, then cooled to -10°C and iron (III) chloride (655mg) added. The reaction was allowed to warm to room temperature overnight,

hydrochloric acid 1N was added and the resulting mixture stirred for one hour. The organic layer was separated, dried (MgSO₄), filtered and the solvent removed under reduced pressure to afford a red oil. The recovered material was dissolved in dry methanol and treated with sodium borohydride (150mg) for one hour. After this time the solvent was removed under reduced pressure and the residue subjected to flash chromatography (methanol/chloroform) to afford the purified product as a clear oil. Treatment of the recovered product with ethereal hydrogen chloride and re-crystallisation from methyl-t-butyl ether gave the title compound as a white powder (210mg). mp 189-190°C. ¹H NMR (360MHz, DMSO-d₆) 3.52 (1H, dd, J = 6.0, 2.0Hz, CHH-NH), 3.74 (1H, dd, J = 6.0, 1.0Hz, CHH-NH), 3.99 (1H, m, CH-CH-NH), 4.63 (1H, d, J = 8.0Hz, CH-CHH-O-CH₂), 4.67 (1H, d, J = 2.0Hz, PhCH-CH), 4.70 (1H, d, J = 9.0Hz, OCHHAr), 4.75 (1H, d, J = 8.0Hz, CH-CHH-OCH₃), 4.80 (1H, d, J = 9.0Hz, OCHHAr), 6.63 (1H, d, J = 3.0Hz, CH-C-CHPh), 7.0-7.3 (9H, m, ArH), 8.03 (1H, s, CF₃C-CH-CCF₃), 8.09 (2H, s, C-CH-CCF₃); m/z (CI⁺) 466; C₂₅H₂₁NOF₆HCl 1/2 H₂O requires: C, 58.77; H, 4.54; N, 2.74; Found C, 58.51; H, 4.47; N, 2.77%.

EXAMPLE 2

4-Phenyl-1,2,3,4-tetrahydroisoquinoline-3-[3',5'-bis
(trifluoromethylbenzyl)carboxylate

a) 4-Phenyl, 3-carboxy-1,2,3,4-tetrahydroisoquinoline

A solution of diphenylalanine (10.0g), concentrated hydrochloric acid (75ml) and formaldehyde (23ml x 37%) was warmed to reflux for 30 minutes. A further quantity of

concentrated hydrochloric acid (20ml) and formaldehyde (20ml x 37%) was added and heating continued for three hours. The solution was then allowed to cool to room temperature overnight. The product was isolated by filtration. Recrystallisation from ethanol afforded the pure product as white needles mp 160-162°C. ¹H NMR (360MHz, DMSO-d₆) 4.54 (1H, m, CHCO₂H), 4.6-4.86 (3H, m, 1H CH-Ph and 2H CH₂-NH), 8.66 (1H, d, J = 6.1Hz, Ar-CH-CH-CHCO₂H). m/z (CI⁺) 253.

10 b) 4-Phenyl-1,2,3,4-tetrahydroisoquinoline-3-[3',5'-bis(trifluoromethyl)benzyl]carboxylate

Di-t-butylidicarbonate (370mg) was added to a rapidly stirring suspension of 4-phenyl-3-carboxy-1,2,3,4-tetrahydroisoquinoline (500mg) and K₂CO₃ (700mg) in a mixture of 1,4-dioxan and water (1:1). The resulting solution was stirred for 4 hours, at room temperature, then acidified with citric acid. The reaction mixture was partitioned between water and ethyl acetate. The organic layers were separated, dried (MgSO₄), filtered and reduced to dryness. The solid residue was re-suspended in dry methanol and caesium carbonate (211mg) added. The resulting solution was stirred for 30 minutes and then the solvent removed under reduced pressure. The solid residue was re-suspended in dry dimethyl formamide (50ml) and a solution of 3,5-bis(trifluoromethyl)benzyl bromide (240µl) added. The resulting solution was stirred at room temperature overnight. The reaction mixture was poured into water (300ml) and extracted into ethyl acetate. The organic extracts were dried (MgSO₄), filtered and the solvent removed under reduced pressure. The oily residue was treated overnight with a saturated solution of hydrochloric acid in methanol. Removal of solvent under reduced pressure and re-crystallisation from

ethanol afforded the product as a white powder. mp 175-176°C. ¹H NMR (360MHz, DMSO-d6) 4.40 (1H, d, J = 10Hz, Ph-CH₂H-N), 4.42 (1H, d, J = 8.0Hz, HN-CH₂-CO₂), 4.61 (1H, d, J = 10Hz, Ph-CH₂H-N), 4.72 (1H, d, J = 8.0Hz, Ph-CH₂-CO₂H), 5.2 (2H, m, CO₂CH₂-Ar), 6.6 (1H, d, J = 5.0Hz, Ar-H-CHPh), 7.1-7.25 (10H, m, ArH), 7.8 (2H, s, 2 x C-CH₂-CCF₃), 8.09 (1H, s, CF₃C-CH₂-CCF₃); m/z (CI⁺) 479; C₂₅H₂₀NO₂F₆Cl requires C, 57.70; H, 3.97; N, 2.69. Found C, 57.70; H, 4.01; N, 2.64%.

EXAMPLE 3

trans-N-Methyl-4-phenyl-3-[3',5'-bis(trifluoromethyl)benzyl]oxy-1,2,3,4-tetrahydroisoquinoline

a) N-Methyl-4-phenyl 3-carboxy-1,2,3,4-tetrahydroisoquinoline

A solution of diphenylalanine (10.0g), concentrated hydrochloric acid (75ml) and formaldehyde (23ml x 37%) was warmed to reflux for 24 hours. The solution was then allowed to cool to room temperature overnight. The product was isolated by filtration. Recrystallisation from ethanol afforded the pure product as white needles mp 174-178°C. ¹H NMR (360MHz, DMSO-d6) 2.19 (3H, s, NCH₃), 4.54 (1H, m, CHCO₂H), 4.6-4.86 (3H, m, 1H CH₂-Ph and 2H CH₂-NH), 8.66 (1H, d, J = 6.1Hz, Ar-CH₂-CH-CHCO₂H). m/z (CI⁺) 267.

b) N-Methyl-4-phenyl-3-hydroxymethyl-1,2,3,4-tetrahydroisoquinoline

To a solution of the N-methyl-4-phenyl, 3-carboxy-1,2,3,4-tetrahydroisoquinoline (2.0g) in dry tetrahydrofuran was added

lithium aluminium hydride (1.0M solution in tetrahydrofuran, 7.1ml). The mixture was heated to reflux for two hours, then allowed to cool to 23°C before being quenched with 4N NaOH. Water (10ml) was added. The mixture was filtered through
 5 celite, the filtrate was extracted with ethyl acetate. The combined organic fractions were dried (MgSO₄), filtered and concentrated under reduced pressure to give the product as a yellow oil (1.49g); ¹H NMR (360MHz, DMSO-d₆), oxalate salt. δ 2.6 (3H, s), 3.2 (1H, m), 3.45 (2H, m), 3.9-4.2 (2H, q), 4.25 (1H,
 10 d), 6.8 (1H, s), 7.73 (8H, m). m/z (E.I.⁺), 222 (M-31); C.I.⁺, 254, 222.

c) N-Methyl-4-phenyl-3-hydroxymethyl-1,2,3,4-tetrahydroisoquinoline (1.0g) and 3,5-bis(trifluoromethyl)
 15 benzylbromide (1.21g) were dissolved in a 1:1 mixture of dry dimethylformamide and tetrahydrofuran (20ml). To this solution was added sodium hydride (60% in oil, 150mg). After two hours, T.L.C. (3:1, hexane:ethyl acetate) shows no starting material. The mixture was poured into water and extracted with
 20 ethyl acetate (2 x 75ml). The combined organic extracts were dried (MgSO₄), filtered and concentrated in vacuo to give a yellow oil. Purified by flash column chromatography (3:1, hexane:ethyl acetate) to give a clear oil. The product was
 25 dissolved in ether, hydrogen chloride gas was bubbled through for two minutes. Ether was removed under reduced pressure to give the title compound as a white crystalline solid (210mg). mp 71°C. NMR, 260MHz, free base CDCl₃. δ 2.5 (3H, s), 3.9 (1H, m), 3.5 (2H, m), 4.2 (1H, d), 4.5 (2H, q), 6.8 (1H, d), 7.74 (8H, m), 7.7 (2H, s), 7.8 (1H, s). m/z, (CI⁺), 480, 481.
 30 (C₂₆H₂₃F₆NO.HCl.1 1/4 H₂O) requires C, 58.00, H, 4.96; N, 2.60 Found C, 58.04; H, 5.33; N, 2.48.

The following examples illustrate pharmaceutical compositions according to the invention.

EXAMPLE 4A Tablets containing 1-25mg of compound

5		<u>Amount mg</u>		
	Compound of formula (I)	1.0	2.0	25.0
	Microcrystalline cellulose	20.0	20.0	20.0
	Modified food corn starch	20.0	20.0	20.0
	Lactose	58.5	57.5	34.5
10	Magnesium Stearate	0.5	0.5	0.5

EXAMPLE 4B Tablets containing 26-100mg of compound

		<u>Amount mg</u>		
	Compound of formula (I)	26.0	50.0	100.0
15	Microcrystalline cellulose	80.0	80.0	80.0
	Modified food corn starch	80.0	80.0	80.0
	Lactose	213.5	189.5	139.5
	Magnesium Stearate	0.5	0.5	0.5
20	The compound of formula (I), cellulose, lactose and a portion of the corn starch are mixed and granulated with 10% corn starch paste. The resulting granulation is sieved, dried and blended with the remainder of the corn starch and the magnesium stearate. The resulting granulation is then compressed into tablets containing			
25	1.0mg, 2.0mg, 25.0mg, 26.0mg, 50.0mg and 100mg of the active compound per tablet.			

EXAMPLE 5 Parenteral injection

		<u>Amount mg</u>
30	Compound of formula (I)	1 to 100mg
	Citric Acid Monohydrate	0.75mg
	Sodium Phosphate	4.5mg
	Sodium Chloride	9mg
	Water for Injections	to 1ml

The sodium phosphate, citric acid monohydrate and sodium chloride are dissolved in a portion of the water. The compound of formula (I) is dissolved or suspended in the solution and made up to volume.

5


EXAMPLE 6 Topical formulation

	<u>Amount mg</u>
Compound of formula (I)	1-10g
Emulsifying Wax	30g
10 Liquid paraffin	20g
White Soft Paraffin	to 100g

The white soft paraffin is heated until molten. The liquid paraffin and emulsifying wax are incorporated and stirred until dissolved. The compound of formula (I) is
15 added and stirring continued until dispersed. The mixture is then cooled until solid.

CLAIMS:

1. A compound of formula (I), or a salt or
5 prodrug thereof:



(I)

wherein:

15 R^1 represents H, C_{1-6} alkyl or phenyl optionally substituted by 1, 2 or 3 groups selected from C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, halo, cyano, nitro, trifluoromethyl, trimethylsilyl, $-OR^a$, SR^a , SOR^a , SO_2R^a , $-NR^aR^b$, $-NR^aCOR^b$, $-NR^aCO_2R^b$, $-CO_2R^a$ and $-CONR^aR^b$;

20 R^2 represents H, C_{1-6} alkyl, COR^7 , $COOR^7$, $CONHR^7$ or phenyl(C_{1-4} alkyl) optionally substituted in the phenyl ring by C_{1-6} alkyl, C_{1-6} alkoxy, halo or trifluoromethyl;

R^3 represents phenyl optionally substituted by 1, 2 or 3 substituents selected from C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, halo, cyano, nitro, trifluoromethyl, trimethylsilyl, $-OR^a$, SR^a , SOR^a , SO_2R^a , $-NR^aR^b$, $-NR^aCOR^b$, $-NR^aCO_2R^b$, $-CO_2R^a$ and $-CONR^aR^b$;

25 each R^4 independently represents C_{1-6} alkyl, C_{1-6} alkoxy, halo or trifluoromethyl;

30 R^5 represents H or C_{1-6} alkyl;

R^6 represents H, C_{1-6} alkyl, or phenyl optionally substituted by 1, 2 or 3 substituents selected from the group consisting of C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, halo, cyano, nitro, trifluoromethyl,

trimethylsilyl, $-OR^a$, SR^a , SOR^a , SO_2R^a , $-NR^aR^b$, $-NR^aCOR^b$,
 $-NR^aCO_2R^b$, $-CO_2R^a$ and $-CONR^aR^b$;

R^7 represents C_{1-6} alkyl or phenyl optionally
substituted by a substituent selected from C_{1-6} alkyl,
5 C_{1-6} alkoxy, halo and trifluoromethyl;

n is 0, 1, 2, 3 or 4;

X represents CH_2 or $C=O$;

Y represents O, S, CH_2 , or NR^{10} where R^{10} is H,
 C_{1-6} alkyl or COR^5 ; or X and Y together represent $CH=CH$;

10 with the proviso that X is not $C=O$ when Y is CH_2 ; and

R^a and R^b each independently represent H,
 C_{1-6} alkyl, trifluoromethyl or phenyl optionally
substituted by C_{1-6} alkyl, halo or trifluoromethyl.

15 2. A compound according to claim 1 wherein R^1
is substituted or unsubstituted phenyl.

20 3. A compound according to claim 1 or claim 2
wherein R^2 is selected from H and C_{1-6} alkyl.

4. A compound according to any preceding
claim wherein R^3 is phenyl substituted by up to three
groups selected from C_{1-6} alkyl, halo and trifluoromethyl.

25 5. A compound according to any preceding
claim wherein Y is selected from O and NR^{10} .

6. A compound according to claim 5 wherein Y
is O.

30 7. A compound according to any preceding
claim wherein R^5 and R^6 each represents H.

8. A compound according to claim 1 selected from:

cis 4-phenyl-3-[3',5'-bis(trifluoromethyl)

benzyloxymethyl] 1,2,3,4-tetrahydroisoquinoline;

5 trans N-methyl-4-phenyl-3-[3',5'-bis(trifluoromethyl)
benzyloxymethyl] 1,2,3,4-tetrahydroisoquinoline;

trans 3',5'-bis(trifluoromethyl)benzyl 4-phenyl 1,2,3,4-
tetrahydroisoquinoline-3-carboxylate;

and salts and prodrugs thereof.

10

9. A pharmaceutical composition comprising a
compound according to any preceding claim in association
with a pharmaceutically acceptable carrier therefor.

15

10. A pharmaceutical composition as claimed in
claim 9 further comprising a bronchodilator.

11. The use of a compound as claimed in claim
1 for the manufacture of a medicament for the treatment
20 of a physiological disorder associated with an excess of
tachykinins.

12. The use of a compound as claimed in claim
1 for the manufacture of a medicament for the treatment
25 of pain or inflammation.

13. The use of a compound as claimed in claim
1 for the manufacture of a medicament for the treatment
or prevention of migraine.

30

14. The use of a compound as claimed in claim
1 for the manufacture of a medicament for the treatment
or prevention of arthritis.

15. The use of a compound as claimed in claim 1 for the manufacture of a medicament for the treatment of postherpetic neuralgia.

-34-

Patents Act 1977
Examiner's report to the Comptroller under
Section 17 (The Search Report)

Application number

GB 9308684.1

Relevant Technical fields

(i) UK Cl (Edition L) C2C

(ii) Int Cl (Edition 5) C07D

Search Examiner

P N DAVEY

Databases (see over)

(i) UK Patent Office

(ii) ONLINE DATABASES: CAS ONLINE

Date of Search

8 JULY 1993

Documents considered relevant following a search in respect of claims

Category (see over)	Identity of document and relevant passages	Relevant to claim(s)
P,X	EP0496369 A1 (HOECHST) 29 July 1992 see example pages 3, 4, 8, 9	1, 3, 5-7
X	EP 0049605 A1 (WARNER-LAMBERT) see example 5, 9, 10	1, 3, 5-7
X	Synth Commun 22(19), 2745-56	1, 3, 5-7
X	Helv Chim Acta 70(7), 1944-54	1, 3, 5-7
X	J Med Chem 29(10) 1953-61	1, 3, 5-7

Category	Identity of document and relevant passages	Relevant to claim(s)

Categories of documents

X: Document indicating lack of novelty or of inventive step.

Y: Document indicating lack of inventive step if combined with one or more other documents of the same category.

A: Document indicating technological background and/or state of the art.

P: Document published on or after the declared priority date but before the filing date of the present application.

E: Patent document published on or after, but with priority date earlier than, the filing date of the present application.

&c: Member of the same patent family, corresponding document.

Databases: The UK Patent Office database comprises classified collections of GB, EP, WO and US patent specifications as outlined periodically in the Official Journal (Patents). The on-line databases considered for search are also listed periodically in the Official Journal (Patents).